




Aspects of Earth Sciences

Prof. Dr. Ulrich A. Glasmacher

Institute of Earth Sciences, Heidelberg University

<u>ulrich.a.glasmacher@geow.uni-heidelberg.de</u>


Representing a group of scientists from HD-Uni, KIT, and Argentine cooperation partner






Why is the Tunnel and the ANDES Underground Laboratory so important for Earth Sciences?


It opens the opportunity to perform Earth Science Research along a **transect** and along **time** in an **Active Orogeny**.









Background: Plate Tectonic Theory









The Andes: An active Orogeny.











Upper Oligocene to Middle Miocene: about 28 – 12 Ma




Upper Permian to lower Triassic: about 260 - 240 Ma
Calculated temperature within the ANDES are about 50 °C





The suggested Earth Sciences Research is two-fold:

- A. During the excavation of the tunnel and the AUL
- **B.** During the Life time of the ANDES Underground Laboratory (AUL).





What do we learn from the Tunnel?

Past Geodynamic movements

by analysing and dating the deformation (structures/faults).

Past Fluid movement and chemistry

by analysing and dating rock alterations (Mineral changes).


Past thermal histories, rock uplift, and erosion

by analysing thermochronometers and applying numerical simulations.

Dynamic of Permian-Triassic and Tertiary volcanic eruptions

by analysing and dating fresh volcanic rocks.

All in Millions of Years





What do we learn from long-term experiments in the ANDES Underground Laboratory


- Recent Geodynamic movements by analysing geophysical parameters.
- Determination of Stress fields and their changes
- Recent Fluid movement and chemistry by analysing fluids.
- Recent Gas movement and chemistry by analysing rock gas.
- Crystallization of melts without influence of Cosmic rays and spallation products.

Continuous Records

Important to mention:

Fluid, Gas data are correlated with geophysical data.

Therefore, monitoring of change of physical parameters is highly important.






A. During the excavation of the tunnel and the ANDES Underground Laboratory.

Suggested Projects:

- Long-term evolution of the Mountain building process using nuclear tracks.
 and a-decay (Glasmacher, Krob, Uni Heidelberg)
- Geochemical alteration in damage zones around tectonic faults.
 (Hilgers, Busch, Kontny, KIT)
- Monitoring of fluid flow and chemistry?
- Quantification and numerical modelling of structural evolution?






A. During the excavation of the tunnel and the ANDES Underground Laboratory.

Suggested Projects:

- Long-term evolution of the Mountain building process using nuclear tracks and a-decay (Glasmacher, Krob, Uni Heidelberg)

Objects: rocks form the tunnel and drill cores





What does "Long-term evolution of the Mountain building process" mean?

Erosion rate [m/Ma]



Rock uplift rate [m/Ma]

Rock uplift rate
> Erosion rate
= Surface elevation
rise



Rock uplift rate
= Erosion rate
= No change
of surface elevation





Rock uplift rate < Erosion rate = Surface elevi

= Surface elevation decrease

ANDES Underground Laboratory











How can this be achieved: **nuclear tracks** and **a-decay**?

- Rocks bear Apatite and Zircon with ²³⁸U and ²³²Th isotopes.
- Apatite and Zircon are archives of time-temperature evolution of rocks/crust.
- Two possible decays: Fissioning and a-decay
- Natural fissioning of ²³⁸U creates linear crystal defects (nuclear tracks) in apatite (AFT) and zircon (ZFT).










How can this be achieved: nuclear tracks and a-decay?

- The linear defects are metastable.
- Increase of temperature erase the tracks (Annealing).
- The annealing process (t-T) is described by equations and t-T evolution numerical modeled.
- a-decay creates ⁴He-isotope in apatite (AHe) and zircon (ZHe).
- The diffusion process (t-T) described by equations and t-T evolution numerical modeled.

Diff. AHe 75°C/1 Ma Anneal. AFT 110°C/10 Ma Diff. ZHe 185°C/1 Ma Anneal. ZFT 330-190°C /10 Ma

Increase of Rock Temperature





Long-term Landscape Evolution


Real Data: Thermochronology and Geol. Evolution

Numerical modelling


HeFTy, QTQt


Pecube, Fetkin,

Petromod















Similar research was performed in the Alps at the Gotthard Tunnel.














B. During the <u>Life time</u> of the ANDES Underground Laboratory (AUL).

Suggested Projects:


- Controlled crystallization of amorphous thin films without cosmic irradiation (Burchard, Glasmacher, Lattard, Uni Heidelberg)
- Long-term monitoring of geogenic gases (Keppler, Glasmacher, Uni Heidelberg)
- Long-term monitoring of magnetic susceptibility of rocks under in-situ stress (Kontny, KIT)
- Long-term rheological and geochemical investigation of host rock integrity in damage zones (Hilgers, Busch, KIT)
- Long-term monitoring of chemical composition of geogene fluids?





Long-term monitoring of geogenic gases (CO₂, N₂, He, H₂ etc. (Keppler, Glasmacher, Uni Heidelberg)


- One small room with open rock surface for the measuring equipment.
 Preferred in zones of damaged rocks
- One small room for computer, power supply.





Long-term monitoring of magnetic susceptibility of rocks under in-situ stress (Kontny, KIT)


- Open rock surface. Preferred in zones of damaged rocks





Long-term rheological and geochemical investigation of host rock integrity in damage zones (Hilgers, Busch, KIT)

- Open rock surface. Preferred in zones of damaged rocks






Controlled crystallization of amorphous thin films without cosmic irradiation (Burchard, Glasmacher, Lattard, Uni Heidelberg)


Experiments performed in the past


- Create a amorphous thin film using an Eximer-laser.
- Analyse by different techniques: Truly amorphous, no nano-crystals.
- Stored for a certain time.
- Fast reheat the amorphous thin film to start slow crystallization (annealing).
- Analyse by different techniques: Shape and distribution of crystals.





Theory: Ideal crystallization of amorphous materials







The crystal growth is described by: Johnson-Mehl-Avrami-Kolmogorov equation (JMAK)

simple form:

$$f(t) = 1 - exp(-At^n)$$
:

A = kinetic crystallization information **| important material constant**|


t = time

n = process constant = q d + B

B = type of seed formation


q = function representing the linear or nonlinear character of the process


d = dimension of seed growth





JMAK parameter A (B q d = constant)





amorphous

PLD layer



Characterization of 2D crystallization

By atomic force microscopy (AFM), difference interference contrast microscopy (DIC), scanning electron microscopy (SEM), fourier transform IR spectroscopy (FTIR)

Thin film on (111) Si-wafer (substrate)

top view

lateral view

uncovered reference

before annealing

amorphous PLD layer

uncovered reference

Silicon-wafer

before annealing

Thin film thickness: ~150 nm


after annealing

(partly crystalline) PLD layer

uncovered reference

Silicon-wafer


Oxidation layer (SiO₂) on wafer surface







Characterization of 2D crystallization

Atomic force microscopy (AFM)












Characterization of 2D crystallization

Difference interference contrast microscopy (DIC)











Characterization of 2D crystallization

Fourier transform IR spectroscopy (FTIR)







Simple time dependent experiments?


Crystallizing thin films (d = 2) of homogenous amorphous material at the same controlled experimental conditions but with variable time should result in simple to evaluable structures.

Based on these data parameter A can easily be calculated.

BUT

All observed experimental parameters are constant.


Crystallization structures is significant different.






The expected


Fo177





Fo178

Slightly different Forsterite thin both crystallized 144h at 1060K.






The unexpected

Fo138






Fo139

Extremely different Forsterite thin films both crystallized 8h at 1098K. Fo139 is completely recrystallized.

Fo139 was stored several month before used for the experiment.





The Results of the Past Experiments

The crystallization structures are statistically related to:


- the storage time of the thin amorphous films before crystallization experiment.
- and not associated with the H₂O / CO₂ contend. (quantified by IR-spectroscopy).

The cause?

The crystallization structure of the same material is extremely sensitive to the formation of crystal seeds.

Hypothesis: Crystal seeds are related to natural high energy radiation.

Therefore, the different crystallization behavior is to be expected directly connected to the flux of natural high energy radiation.






Conclusion for the ANDES experiments

Experiments with samples produced and processed under low radiation flux conditions are necessary to gain the "real" material crystallization constants.

The Results of these experiments will be compared with results from crystallization runs performed under natural flux conditions





Controlled crystallization of amorphous thin films without cosmic irradiation (Burchard, Glasmacher, Lattard, Uni Heidelberg)


- An area within the AUL to perform the experiments.
- Equipment:

Eximer-Laser, Power etc.: to generate the thin films

Storage area

Oven: performing the annealing experiments

Analyser: First analysing of crystallization structure





B. During the **Life time** of the ANDES Underground Laboratory (AUL).

Suggested Projects:

- Controlled crystallization of amorphous thin films without cosmic irradiation (Burchard, Glasmacher, Lattard, Uni Heidelberg)
- Long-term monitoring of geogenic gases (Keppler, Glasmacher, Uni Heidelberg)
- Long-term monitoring of magnetic susceptibility of rocks under in-situ stress (Kontny, KIT)
- Long-term rheological and geochemical investigation of host rock integrity in damage zones (Hilgers, Busch, KIT)
- Long-term monitoring of chemical composition of geogene fluids?

Muchas Gracias!

Thank you very much!